

UC Irvine

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health

Title

Emergency Medicine Research Priorities for Early Intervention for Substance Use Disorders

Permalink

<https://escholarship.org/uc/item/2806d7kc>

Journal

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health, 20(2)

ISSN

1936-900X

Authors

Hawk, Kathryn F.
Glick, Rachel L.
Jey, Aurthur R.
[et al.](#)

Publication Date

2019

DOI

10.5811/westjem.2019.1.39261

Supplemental Material

<https://escholarship.org/uc/item/2806d7kc#supplemental>

License

[CC BY 4.0](#)

Peer reviewed

Emergency Medicine Research Priorities for Early Intervention for Substance Use Disorders

Kathryn F. Hawk, MD, MHS*
Rachel L. Glick, MD†
Arthur R. Jey, MD‡
Sydney Gaylor, BS§
Jamie Doucet, MPH¶
Michael P. Wilson, MD, PhD||
John S. Rozel, MD#

*Yale University, Department of Emergency Medicine, New Haven, Connecticut
†University of Michigan, Department of Psychiatry, Ann Arbor, Michigan
‡Sutter Medical Center, Department of Emergency Medicine, Sacramento, California
§University of California- San Diego, Department of Emergency Medicine, San Diego, California
¶Rosalind Franklin University, Chicago, Illinois
||University of Arkansas for Medical Sciences, Department of Emergency Medicine, Little Rock, Arkansas
#University of Pittsburgh, Department of Psychiatry, Pittsburgh, Pennsylvania

Section Editor: Marc L. Martel, MD

Submission history: Submitted June 4, 2018; Revision received January 22, 2019; Accepted January 25, 2019

Electronically published February 19, 2019

Full text available through open access at http://escholarship.org/uc/uciem_westjem

DOI: 10.5811/westjem.2019.1.39261

Introduction: Patients with substance use disorders (SUDs) frequently seek emergency care, and the emergency department (ED) may be their only point of contact with the healthcare system. While the ED visit has been increasingly recognized as providing opportunity for interventions around substance use, many questions remain.

Methods: In December 2016 the Coalition on Psychiatric Emergencies (CPE) convened the first Research Consensus Conference on Acute Mental Illness, which consisted of clinical researchers, clinicians from emergency medicine, emergency psychiatry, emergency psychology, representatives from governmental agencies and patient advocacy groups. Background literature review was conducted prior to the meeting, and questions were iteratively focused, revised, voted on and ranked by perceived importance using nominal group method.

Results: The main goal of the SUD workgroup was to identify research priorities and develop a research agenda to improve the early identification of and management of emergency department (ED) patients with SUDs with the goal of improving outcomes. This article is the product of a breakout session on “Special Populations: Substance Use Disorder.” The workgroup identified with high consensus six research priorities for their importance related to the care of ED patients with SUDs in these overall domains: screening; ED interventions; the role of peer navigators; initiation of SUD management in the ED; specific patient populations that may impact the effectiveness of interventions including sociogenerational and cultural factors; and the management of the acutely intoxicated patient.

Conclusion: Emergency providers are increasingly recognizing the important role of the ED in reducing adverse outcomes associated with untreated SUDs. Additional research is required to close identified knowledge gaps and improve care of ED patients with SUD. [West J Emerg Med. 2019;20(2)386–392.]

INTRODUCTION

In 2016, the National Survey on Drug Use and Health (NSDUH) demonstrated that only 2.3 million of the 20.5 million individuals with an identified need for treatment of a

substance use disorder (SUD), had received care within the prior year.¹ Nonetheless, patients with SUD frequently seek emergency care, making up half of the more than 4.9 million emergency department (ED) visits for drug-related complaints.²

Patients with unmet treatment needs are more likely to be hospitalized than those receiving treatment for a SUD, and substance use is associated with higher rates of unintentional injuries, motor vehicle collisions, interpersonal violence, human immunodeficiency virus infection, and intentional or accidental overdose.³⁻⁶ Increasingly, the ED visit has been identified as a unique opportunity for intervention and linkage to treatment for patients who are at risk for or who currently have SUDs, whether for tobacco, opioid or alcohol use.⁷⁻¹⁰

The ED may be the only point of contact with the healthcare system for some patients with SUDs. An ED visit for an acute injury, illness or overdose may provide a window of opportunity where patients are more receptive to education about and referral to treatment for SUD.¹¹ Over the past decade, significant strides have been made in the field of ED-based identification, interventions and referrals for the treatment of SUD, but many questions remain. The goal of the SUD workgroup of the Coalition on Psychiatric Emergencies Research Consensus Conference on Acute Mental Illness was to identify research priorities and develop a research agenda to improve the early identification of and management of ED patients with SUDs with the goal of improving outcomes.

METHODS

Please see the Executive Summary (Appendix) for full methods. Participants from a variety of disciplines – emergency medicine (EM), emergency psychiatry, emergency psychology, clinical research, governmental agencies, and patient advocacy groups – were invited to participate in a research consensus session held prior to a joint emergency-psychiatry conference (the 7th Annual National Update on Behavioral Emergencies). Background literature reviews were performed prior to the in-person meeting. A total of 38 articles were circulated to the SUD group in advance. The working group initially identified three key areas: identification and diagnosis/screening; ED-based interventions; and linkages to the continuum of care. During the conference, the group spent time discussing research gaps related to addiction in the ED and identified 36 research topics. After spending time generating the initial 36 questions, the group re-reviewed them and held additional discussions to add clarity and intent; it then condensed the list to 24 questions.

A nominal group technique was employed to develop group consensus on the highest priority research gaps. Each member was given five points with which to vote for the questions they felt were most important. The questions were then ranked by the number of votes. The group identified six key research questions to guide ED-based interventions for SUD. Following the nominal technique, additional input was solicited from participants, questions were iteratively focused and revised, voted on, and then ranked by importance. Following the in-person session, the workgroup developed additional consensus by meeting electronically to further refine the final form of each question.

The working group focused on SUD was made up of seven people: one EM clinician researcher, one EM clinician, and two clinician psychiatrists; a non-physician student; a participant from a medical association, and an observer from industry. The average age of the participants was about 42 years old; four were females and three were males.

RESULTS

During the consensus conference, research questions and topics were sequentially proposed by individual members of the workgroup, and were transcribed into a large working board visible to all group members in real time. Workgroup members proposed research topics individually and in a sequential fashion, for a total of 36 research topics. Topics were discussed, grouped, voted on, and prioritized using the nominal group technique.

DISCUSSION

The workgroup identified six questions as the highest priority areas for early identification of SUDs in the ED. (Additional questions and discussion, organized by topic, are also included in Table 1 and Table 2.)

Screening

Based on a robust literature search, the United States Preventive Services Task Force (USPSTF) and the American College of Emergency Physicians (ACEP) recommend screening and brief intervention for alcohol use disorders in primary care and ED settings.¹¹⁻¹³ The role of universal screening for illicit drug use, either in the primary care or ED setting, is less clear. Although evidence is lacking, increased rates of illicit drug use among ED patients, recent increases in opioid-associated mortality, and recent ED-based studies showing improved outcomes after ED intervention, provide a basis for the role of ED screening for SUDs.^{8,12,14-16}

While ED-based research studies focused on SUDs have used screening to identify potential study subjects, little is known about either the impact or the most effective implementation of ED-wide screening procedures in the day-to-day functioning of an ED. Multiple studies have adapted, developed and piloted a variety of screening tools for SUDs using tablet- and kiosk-based platforms, but consensus on the most effective implementation of ED-based screening algorithms outside of a research study has not been reached.^{2,7,17,18} Several EDs have implemented Screening, Brief Intervention and Referral Treatment (SBIRT) programs including use of health promotion advocates such as in Project Alcohol and Substance Abuse Services, Education, and Referral to Treatment (ASSERT), or training ED residents and faculty as part of Substance Abuse and Mental Health Services Administration's (SAMSHA) SBIRT training grants as best practice.¹⁹

Table 1. Key research questions to guide emergency department-based interventions for substance use disorders.

Question 1	What are the most effective, efficient and appropriate ways to screen for SUD in the ED?
Question 2	What are the most effective ED-based interventions for SUD?
Question 3	What is the role for initiation and management of SUD treatment and detoxification in the ED?
Question 4	What is the role of sociocultural and generational factors in acceptability, accessibility, and benefit of ED-based initiatives?
Question 5	What are the best practices for the evaluation and management of the acutely intoxicated patient?
Question 6	What role can peer mentors, or patient navigators, play in improving patient outcomes?

SUD, substance use disorder; *ED*, emergency department.

Table 2. Key research questions to guide efforts for improved care of individuals with substance use disorders.

Topic area 1: What are the most effective, efficient and appropriate ways to screen for SUD in the ED?
What is the best approach for sensitively and effectively screening for drug and alcohol use in the ED?
How effective are current screening tools in different populations and do results vary with patient characteristics/identity: generational (i.e., millennial vs geriatric), gender, religious, cultural factors?
What is the most cost-effective way to implement high-quality ED-based screening for SUD ?
What is the role for SUD screening in ED triage? For universal screening?
Topic area 2: What are the most effective ED-based interventions for SUD?
Which ED-based interventions can reduce cost, reduce mortality and increase treatment adherence?
Do harm reduction initiatives (i.e., overdose prevention education, naloxone distribution) improve outcomes?
To which types of treatment/services should ED patients with SUD be referred?
Topic area 3: What is the role for initiation and management of SUD treatment and detoxification in the ED?
Who is appropriate for ED-initiated outpatient treatment of alcohol withdrawal?
Is there a need for development of a validated ED-based protocol for initiating outpatient treatment of alcohol withdrawal?
Is there a need for development of a validated ED-based protocol for initiating buprenorphine for the treatment of OUD, including who is most likely to benefit and when?
Topic area 4: What is the role of sociocultural and generational factors in acceptability, accessibility, and benefit if ED-based initiatives?
Topic area 5: What are the best practices for the evaluation and management of the acutely intoxicated patient?
Do better evaluation, diagnosis and treatment of agitation of patients with acute intoxication of traditional drugs of abuse as well as newly emerging novel psychoactive substances (NPS) currently exist?
Are there clinical guidelines for management of acute stimulant intoxication?
What is the appropriate role of drug/toxicology screens in the ED?
Is there evidence based-criteria for medical workup prior to psychiatric evaluation?
Topic area 6: What role can peer mentors, or patient navigators, play in improving patient outcomes?
What role can peer mentors, or patient navigators, play in improving patient outcomes?

SUD, substance use disorder; *ED*, emergency department; *NPS*, novel psychoactive substances; *OUD*, opioid use disorder.

Although several of these programs are of long standing and have linked thousands of ED patients to SUD care, the most effective and efficient way to screen in diverse ED settings remains unclear.¹² Importantly, the logistics of who administers the screen and how it is performed (e.g., triage nurse; tablet-based or self-administered; emergency provider) will influence the overall acceptability of the process to the patient, the sensitivity to detect SUDs, the integration of the

process into the ED workflow, and the overall sustainability. The most efficient and effective approach to screening for SUDs will likely vary based on patient population, geography, ED volume, community resources, ED staffing and academic vs community hospital settings, and may vary across cultural and generational patients within ED populations. Increasingly, the ED has been recognized as an important venue to identify and engage patients with SUDs.¹³

Intervention

Although significant strides have been made in improving outcomes for ED patients with risky alcohol use, uncertainty surrounding the most effective interventions to reduce illicit drug use persist. A fairly robust literature exists supporting the implementation of SBIRT for alcohol use disorders in primary care settings^{11,20} although mixed results have been seen in the ED.^{7,21,22} Brief interventions incorporate principles of motivational interviewing, an evidence-based counseling technique that uses empathy, positive framing, reflective listening, and gentle education to enhance motivation to reduce risky behaviors.^{10,23} Brief interventions for patients with at-risk or hazardous drinking usually focus on reducing use, while the focus for patients with dependence is on enhancing motivation to accept a referral to formal treatment.¹³⁻¹⁵ Some ED-based studies have shown success in reducing alcohol consumption, episodes of binge drinking and episodes of driving after drinking in harmful and hazardous alcohol drinkers, although other studies have been less encouraging, with no persistent effect at one year.^{7,22,24,25}

ED-based brief interventions for drug use have been less promising. The Screening, Motivational Assessment, Referral, and Treatment in Emergency Departments (SMART-ED) Clinical Trials Network Study across six academic EDs did not detect differences in drug use at any point in time.¹⁶ Additionally, a single, large, randomized control trial (RCT) found that a brief motivational intervention for patients with alcohol or drug use disorders did not improve attendance at post-ED intervention over a case management intervention.^{16,18,19} However, there were several methodological issues with these studies, and it is likely that one intervention may not be effective for all types of drugs at all levels of severity.

More recently, ED-based interventions specific to patients with opioid use disorders (OUD) have shown more promise. One pilot RCT of ED patients with non-medical opioid use found a significant reduction in overdose-risk behaviors and a reduction in non-medical opioid use at six months after an ED-based motivational interview intervention compared to usual care.⁹ In one single ED-based RCT, patients with opioid dependence who received a brief intervention and ED-initiation of buprenorphine were significantly more likely to be engaged in treatment for OUD at 30 days (78% vs 37%) and had fewer days of opioid use than the standard referral to treatment group.²⁰ This study, augmented by the persistent rapid rise of opioid-associated fatalities, has prompted a number of EDs across the country to develop programs initiating treatment with buprenorphine for OUD in the ED, although many questions remain about how to optimize implementation, patient selection, models of linkage and induction/dosing algorithms to maximize safe and effective linkage to treatment.^{21,22} Studies are needed to optimize these and other strategies to enhance the success of ED-initiated buprenorphine, and to better characterize patient

and provider facilitators and barriers to the implementation of this intervention.

In the general population in the Western world, approximately 10% of women and 20% of men will have an alcohol use disorder (AUD).²³ About 50% of individuals with AUD are expected to have withdrawal symptoms with reduction or cessation of alcohol use, and 3-5% will have severe complications of withdrawal including seizures or delirium.²³ That said, ED clinicians routinely care for those with the highest risk of complicated withdrawal. General consensus and non-ED based literature suggest patients with mild to moderate AUD may be appropriate for outpatient management with or without oral benzodiazepines.^{29,30} However, there is a paucity of prospective, ED-based studies to provide guidance for the ED population.

Patients at high risk for severe withdrawal and therefore generally inappropriate for outpatient management, include those with a history of alcohol withdrawal seizure or delirium, psychiatric or medical co-morbidities, or patients who receive multiple doses of benzodiazepines without significant reduction in Clinical Institute Withdrawal Assessment scale.²⁴ Clinical decisions regarding the disposition of patients at risk for or with symptoms of alcohol withdrawal are challenging given the dearth of prospective, evidence-based ED studies to guide the risk-benefit analysis of discharging the patient who is at risk for moderate to severe alcohol withdrawal. Moreover, although multiple outpatient regimens for the treatment of alcohol withdrawal symptoms have been described, no clear evidence exists for the most appropriate medication type and dosing schedule.^{23,25,26}

Patient Population Factors

Little is known about the role of sociocultural and generational factors in the acceptability, accessibility, and benefit of ED-based initiatives to reduce harmful substance use and provide linkage to treatment for SUD. Many novel interventions rely on relatively new mobile health and other technology, including smartphone, text messaging and videoconferencing-based interventions, or wearable biosensors, which may be more appealing to younger patients, but create an additional barriers for identifying or intervening in substance use for populations with less intrinsic exposure to technology because of cultural factors or age.²⁷⁻²⁹ Although intervention developers may be specifically targeting younger patients, cultural and generational factors should be considered in the development and implementation of ED-based initiatives given the pervasive distribution of SUDs across all demographics.¹⁻³

Initial Substance Use Disorder Management

Intoxicated patients present unique challenges to the emergency physician. They can be agitated and disruptive.³⁰ Patients present with alcohol intoxication alone or in combination with other drugs, but an increasing number of

visits are due to stimulants, novel psychoactive substances (NPS) and designer drugs.³¹ Literature on management of alcohol intoxication exists but is built on consensus and our limited knowledge of treating agitation in general. Little research has been done on the best management for stimulants and newer substances and the current literature consists mainly of descriptive small series, case reports, or surveys of clinicians' experiences.³

Of the more than 4.5 million ED visits in 2009 for drug-related causes,³⁴ 32% involved alcohol use alone or in combination with other drugs. Nearly 94,000 visits were for stimulants and over 400,000 were for cocaine, while fewer were for phencyclidine, gamma hydroxybutyrate, and ecstasy.³⁵ While it is not clear how many of these visits were for substance-related intoxication as opposed to withdrawal, drug seeking or other reasons, it is clear that the intoxicated patient presents unique challenges to the ED treatment team. Furthermore, the burden of caring for patients with acute, alcohol-related visits more than doubled between 2001 and 2011 reflecting an increased number of visits, longer length of stay, and more intensive use of diagnostic services.³⁶

When a patient presents with suspected drug intoxication and is sleepy or sedated, management is straightforward and supportive until the substance clears and the patient awakens. When a patient is agitated, disruptive, and not cooperative, management is more difficult. Management of agitation in general is not well studied,³⁷ and it is not surprising that our understanding of the best approach to managing the patient who is agitated because of intoxication is limited and based more on retrospective reviews, anecdotal information, and expert consensus.^{38,39} Experts recommend identification of drugs/alcohol as cause of agitation as a first step, followed by verbal de-escalation and medications as necessary, but the research to back this approach is lacking.⁴⁰⁻⁴¹

Another issue to consider when treating intoxicated patients in the emergency setting is the value of laboratory testing such as drug screens and blood alcohol levels. Available toxicology screens often miss substances, and patient history may be more helpful than expensive diagnostic tests except in situations where patients are obtunded or otherwise unable to provide a history.⁴²⁻⁴³ As clinical intoxication frequently does not align with blood alcohol levels, questions frequently arise in determining patient ability to make medical decisions, including the ability to refuse medical care or, depending on the ED setting, when patients are appropriate for psychiatric evaluation.⁴⁴⁻⁴⁶ No clear evidence-based consensus currently exists on the best practices for medical workup prior to psychiatric evaluation.

Substance use is a well-known risk factor for suicide, and a large percentage of individuals who die by suicide are intoxicated at the time of their deaths.⁴⁷ One challenge is how to best assess risk of suicide in the patient who presents to the ED with suicidal statements when intoxicated but later recants when

sober saying they either "just said those things" because they were intoxicated or denying any memory of making suicidal statements or having suicidal thoughts. Persistent knowledge gaps exist around best practices for this ED population.

Peer Mentors

Peer mentors, people with the lived experience of recovery from addiction and mental illness, are becoming increasingly common in the healthcare landscape.⁴⁸ Peer mentors have been identified as a potential bridge to treatment for ED patients after non-fatal opioid overdose, although the impact of this approach on outcomes is unclear.⁴⁹ Early indicators suggested that using peer mentors and peer-led programs can be a helpful diversion for people with addiction and mental health emergencies.⁵⁰⁻⁵¹ Larger studies have shown limited benefit from peer interventions, often due to inconsistent program fidelity and heterogeneous approaches.⁵² Emerging efforts to create fidelity models are promising.⁵³ However, interventions need to be evidenced based and administered by individuals adhering to critical actions with routine fidelity checks and supervision. Additional research to explore the impact of a potentially important and effective way to support and engage people with addiction emergencies, including after opioid overdose, who require linkage to early recovery resources are needed.

LIMITATIONS

There are several limitations to this study. First, this was not a structured review of literature but rather the outcome of an expert consensus group meeting that was held in 2016. By the time of this paper's publication, it is possible that studies may have been conducted that answer or speak to some of the highlighted questions raised. Second, the group focus was narrowed to the early identification and management of patients presenting to the ED with SUDs, drugs and alcohol. Although we recognize the impact on tobacco use disorder and other medical and psychiatric comorbidities, given our limited time, we limited the scope of our work to the care and management of SUDs in the acute care settings and thus we did not specifically discuss tobacco or include focus on the management of other comorbidities. As with many in-person consensus conferences, participation is limited to those who were able to travel and attend in person; had all of the original invitees or others with valuable experience been able to attend, the findings may have been different. Nonetheless, we have highlighted a number of priority areas in which additional research is clearly needed and that can guide ongoing research as we work to improve outcomes of ED patients with SUDs.

CONCLUSION

Emergency providers are increasingly recognizing the important public health role that EDs can play to reduce adverse outcomes associate with undiagnosed and untreated

substance use disorder. Much like the ED is the “front door” for hospital admission, it is also a portal to the community to identify patients with SUDs and to deliver interventions to improve outcomes.

ACKNOWLEDGMENTS

The authors would like to acknowledge the Clinician Scientist Program at the University of Arkansas for Medical Sciences, which partially supported Dr. Wilson during this work. The authors would also like to thank Gail D’Onofrio for her review and feedback, and Loren Rives for her support in preparation of this manuscript.

Address for Correspondence: Kathryn Hawk, MD, Yale University, Department of Emergency Medicine, 464 Congress Ave, Suite 260, New Haven CT 06519. Email: kathryn.hawk@yale.edu.

Conflicts of Interest: By the WestJEM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. The first author (KH) was partially supported by NIDA Grant 4K12DA033312-04 during this work. The Coalition is funded, in part, by an unrestricted grant from Alexza.

Copyright: © 2019 Hawk et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

- Ahmsbrak R, Bose J, Hedden SL, et al. Key substance use and mental health indicators in the United States: Results from the 2016 National Survey on Drug Use and Health. *SAMHSA*. 2017.
- The DAWN Report: Highlights of the 2010 Drug Abuse Warning Network (DAWN) findings on drug-related emergency department visits. Available at: <http://archive.samhsa.gov/data/2k12/DAWN096/SR096EDHighlights2010.htm>. Accessed April 19, 2015.
- Konstantopoulos WLM, Dreifuss JA, Mcdermott KA, et al. Identifying patients with problematic drug use in the emergency department: results of a multisite study. *Ann Emerg Med*. 2014;64(5):516-25.
- Rockett IRH, Putnam SL, Jia H, et al. Unmet substance abuse treatment need, health services utilization, and cost: A population-based emergency department study. *Ann Emerg Med*. 2005;45(2):118-27.
- Hall AJ, Logan JE, Toblin RL, et al. Patterns of abuse among unintentional pharmaceutical overdose fatalities. *JAMA*. 2008;300(22):2613-20.
- Florence CS, Zhou C, Luo F, et al. The Economic Burden of Prescription Opioid Overdose, Abuse, and Dependence in the United States, 2013. *Med Care*. 2016;54(10):901-6.
- Bernstein SL, D’Onofrio G, Rosner J, et al. Successful tobacco dependence treatment in low-income emergency department patients: a randomized trial. *Ann Emerg Med*. 2015;66(2):140-7.
- D’Onofrio G, Fiellin DA, Pantalon MV, et al. A brief intervention reduces hazardous and harmful drinking in emergency department patients. *Ann Emerg Med*. 2012;60(2):181-92.
- Bohnert ASB, Bonar EE, Cunningham R, et al. A pilot randomized clinical trial of an intervention to reduce overdose risk behaviors among emergency department patients at risk for prescription opioid overdose. *Drug Alcohol Depend*. 2016;163:40-7.
- D’Onofrio G, O’Connor PG, Pantalon MV, et al. Emergency department-initiated buprenorphine/naloxone treatment for opioid dependence: a randomized clinical trial. *JAMA*. 2015;313(16):1636-44.
- Bernstein SL, D’Onofrio G. A promising approach for emergency departments to care for patients with substance use and behavioral disorders. *Health Aff (Millwood)*. 2013;32(12):2122-8.
- D’Onofrio G, Degutis LC. Integrating Project ASSERT: A screening, intervention, and referral to treatment program for unhealthy alcohol and drug use into an urban emergency department. *Acad Emerg Med*. 2010;17(8):903-11.
- Hawk K, D’Onofrio G. Emergency department screening and interventions for substance use disorders. *Addict Sci Clin Pract*. 2018;13(1):18.
- D’Onofrio G, Pantalon MV, Degutis LC, et al. Brief intervention for hazardous and harmful drinkers in the emergency department. *Ann Emerg Med*. 2008;51(6):742-50.e2.
- Glass JE, Andréasson S, Bradley KA, et al. Rethinking alcohol interventions in health care: a thematic meeting of the International Network on Brief Interventions for Alcohol & Other Drugs (INEBRIA). *Addict Sci Clin Pract*. 2017;12(1):14.
- Bogenschutz MP, Donovan DM, Mandler RN, et al. Brief intervention for patients with problematic drug use presenting in emergency departments: a randomized clinical trial. *JAMA Intern Med*. 2014;174(11):1736-45.
- Donovan DM, Bogenschutz MP, Perl H, et al. Study design to examine the potential role of assessment reactivity in the Screening, Motivational Assessment, Referral, and Treatment in Emergency Departments (SMART-ED) protocol. *Addict Sci Clin Pract*. 2012;7(1):16.
- Blow FC, Walton MA, Murray R, et al. Intervention attendance among emergency department patients with alcohol-and drug-use disorders. *J Stud Alcohol Drugs*. 2010;71(5):713-9.
- Young MM, Stevens A, Galipeau J, et al. Effectiveness of brief interventions as part of the Screening, Brief Intervention and Referral to Treatment (SBIRT) model for reducing the nonmedical use of psychoactive substances: a systematic review. *Syst Rev*. 2014;3:50.
- D’Onofrio G, O’Connor PG, Pantalon MV, et al. Emergency department-initiated buprenorphine/naloxone treatment for opioid dependence: a randomized clinical trial. *JAMA*. 2015;313(16):1636-44.
- Rubin R. As overdoses climb, emergency departments begin treating opioid use disorder. *JAMA*. 2018;319(21):2158-60.

22. Duber HC, Barata IA, Cioè-Peña E, et al. Identification, management, and transition of care for patients with opioid use disorder in the emergency department. *Ann Emerg Med.* 2018;72(4):420–31.
23. Longo DL, Schuckit MA. Recognition and management of withdrawal delirium (delirium tremens). *N Engl J Med.* 2014;371(22):2109–13.
24. Yanta JH, Swartzentruber GS, Pizon AF. Alcohol withdrawal syndrome: improving outcomes through early identification and aggressive treatment strategies. *Emerg Med Pract.* 2015;17(6):1–18; quiz 19.
25. Hendey GW, Dery RA, Barnes RL, et al. A prospective, randomized, trial of phenobarbital versus benzodiazepines for acute alcohol withdrawal. *Am J Emerg Med.* 2011;29(4):382–5.
26. Askgaard G, Hallas J, Fink-Jensen A, et al. Phenobarbital compared to benzodiazepines in alcohol withdrawal treatment: A register-based cohort study of subsequent benzodiazepine use, alcohol recidivism and mortality. *Drug Alcohol Depend.* 2016;161:258–64.
27. Suffoletto B, Kristan J, Callaway C, et al. A text message alcohol intervention for young adult emergency department patients: a randomized clinical trial. *Ann Emerg Med.* 2014;64(6):664–72.e4.
28. Panagioti M, Reeves D, Meacock R, et al. Is telephone health coaching a useful population health strategy for supporting older people with multimorbidity? An evaluation of reach, effectiveness and cost-effectiveness using a 'trial within a cohort.' *BMC Med.* 2018;16(1):80.
29. Carreiro S, Smelson D, Ranney M, et al. Real-time mobile detection of drug use with wearable biosensors: a pilot study. *J Med Toxicol.* 2015;11(1):73–9.
30. Thomas J, Moore G. Medical-legal Issues in the agitated patient: cases and caveats. *West J Emerg Med.* 2013;14(5):559-65.
31. Schifano F, Chiappini S, Corkery JM, et al. Abuse of prescription drugs in the context of novel psychoactive substances (NPS): A Systematic Review. *Brain Sci.* 2018;8(4):73.
32. Bui QM, Simpson S, Nordstrom K. Psychiatric and medical management of marijuana intoxication in the emergency department. *West J Emerg Med.* 2015;16(3):414–7.
33. Vazirian M, Jerry JM, James J, et al. Bath salts in the emergency department. *J Addict Med.* 2015;9(2):94–8.
34. Center for Health Statistics N. National Hospital Ambulatory Medical Care Survey: 2010 Emergency Department Summary Tables 2010. Available at: https://www.cdc.gov/nchs/data/ahcd/nhamcs_emergency/2010_ed_web_tables.pdf. Accessed December 15, 2016.
35. DrugFacts: Drug-related hospital emergency room visits | National Institute on Drug Abuse (NIDA). Available at: <http://www.drugabuse.gov/publications/drugfacts/drug-related-hospital-emergency-room-visits>. Accessed September 29, 2015.
36. Mullins PM, Mazer-Amirshahi M, Pines JM. Alcohol-related Visits to US emergency departments, 2001–2011. *Alcohol Alcohol.* 2017;52(1):119–25.
37. Holloman GH, Zeller SL. Overview of Project BETA: Best practices in evaluation and treatment of agitation. *West J Emerg Med.* 2012;13(1):1–2.
38. MacDonald K, Wilson MP, Minassian A, et al. A retrospective analysis of intramuscular haloperidol and intramuscular olanzapine in the treatment of agitation in drug- and alcohol-using patients. *Gen Hosp Psychiatry.* 2010;32(4):443–5.
39. D'Onofrio G, Jauch E, Jagoda A, et al. NIH Roundtable on Opportunities to Advance Research on Neurologic and Psychiatric Emergencies. *Ann Emerg Med.* 2010;56(5):551–64.
40. Stowell K, Florence P, Harman H, et al. Psychiatric Evaluation of the Agitated Patient: consensus statement of the American Association for Emergency Psychiatry Project BETA Psychiatric Evaluation Workgroup. *West J Emerg Med.* 2012;13(1):11–6.
41. Wilson MP, Pepper D, Currier GW, et al. The psychopharmacology of agitation: consensus statement of the American Association for Emergency Psychiatry Project Beta psychopharmacology workgroup. *West J Emerg Med.* 2012;13(1):26–34.
42. Bhalla A. Bedside point of care toxicology screens in the ED: utility and pitfalls. *Int J Crit Illn Inj Sci.* 2014;4(3):257–60.
43. Tomassoni AJ, Hawk KF, Jubanyik K, et al. Multiple fentanyl overdoses — New Haven, Connecticut, June 23, 2016. *MMWR Morb Mortal Wkly Rep.* 2017;66(4):107–11.
44. Volz TM, Boyer KS. The development of a behaviorally-based alcohol intoxication scale. *J Emerg Nurs.* 2014;40(4):330–5.
45. Olson KN, Smith SW, Kloss JS, et al. Relationship between blood alcohol concentration and observable symptoms of intoxication in patients presenting to an emergency department. *Alcohol Alcohol.* 2013;48(4):386–9.
46. McCormack RP, Gallagher T, Goldfrank LR, et al. Including frequent emergency department users with severe alcohol use disorders in research: assessing capacity. *Ann Emerg Med.* 2015;65(2):172–7.e1.
47. Norström T, Rossow I. Alcohol Consumption as a risk factor for suicidal behavior: a systematic review of associations at the individual and at the population level. *Arch Suicide Res.* 2016;20(4):489–506.
48. Myrick K, del Vecchio P. Peer support services in the behavioral healthcare workforce: State of the field. *Psychiatr Rehabil J.* 2016;39(3):197-203.
49. Samuels E. Emergency department naloxone distribution: a Rhode Island Department of Health, recovery community, and emergency department partnership to reduce opioid overdose deaths. *R I Med J.* 2014;97(10):38–9.
50. Shattell MM, Harris B, Beavers J, et al. A recovery-oriented alternative to hospital emergency departments for persons in emotional distress: "The Living Room." *Issues Ment Health Nurs.* 2014;35:4–12.
51. Parham K, Rozel JS. Peer professionals benefit psychiatric care. *Psychiatry Advisor.* Available at: <http://www.psychiatryadvisor.com/opinion/peer-professionals-benefit-psychiatric-care/article/368458/>. Accessed March 20, 2017.
52. Lloyd-Evans B, Mayo-Wilson E, Harrison B, et al. A systematic review and meta-analysis of randomised controlled trials of peer support for people with severe mental illness. *BMC Psychiatry.* 2014;14:39.
53. Chinman M, Mccarthy S, Mitchell-Miland C, et al. Early stages of development of a peer specialist fidelity measure. *Psychiatr Rehabil J.* 2016;39(3):256-65.